
2007 Hypatia Contest (Grade 11) Wednesday, April 18, 2007

1. The diagram shows four cities A, B, C, and D, with the distances between them in kilometres.



- (a) Penny must travel from A through each of the other cities exactly once and then back to A. An example of her route might be $A \to B \to D \to C \to A$. List all routes that Penny could travel.
- (b) Identify one route of the shortest possible length and one of the longest possible length. Explain how you obtained your answer.
- (c) Just before leaving A, Penny learns that
 - she must visit a fifth city E,
 - ullet E is connected directly to each of A, B, C, and D, and
 - E must be the third city she visits.

Therefore, the trip would be $A \to \underline{\hspace{1cm}} \to \underline{\hspace{1cm}} \to E \to \underline{\hspace{1cm}} \to A$.

How many different routes are now possible? Explain how you obtained your answer.

(d) The trip $A \to D \to C \to E \to B \to A$ is 600 km long.


The trip $A \to C \to D \to E \to B \to A$ is 700 km long.

The distance from D to E is 225 km.

What is the distance from C to E? Explain how you obtained your answer.

- 2. Olayuk has four pails labelled P, Q, R, and S, each containing some marbles. A "legal move" is to take one marble from each of three of the pails and put the marbles into the fourth pail.
 - (a) Initially, the pails contain 9, 9, 1, and 5 marbles. Describe a sequence of legal moves that results in 6 marbles in each pail.
 - (b) Suppose that the pails initially contain 31, 27, 27, and 7 marbles. After a number of legal moves, each pail contains the same number of marbles.
 - i. Describe a sequence of legal moves to obtain the same number of marbles in each pail.
 - ii. Explain why at least 8 legal moves are needed to obtain the same number of marbles in each pail.
 - (c) Beginning again, the pails contain 10, 8, 11, and 7 marbles. Explain why there is no sequence of legal moves that results in an equal number of marbles in each pail.

- 3. Consider the quadratic function $f(x) = x^2 4x 21$.
 - (a) Determine all values of x for which f(x) = 0 (that is, $x^2 4x 21 = 0$).
 - (b) If s and t are different real numbers such that $s^2 4s 21 = t^2 4t 21$ (that is, f(s) = f(t)), determine the possible values of s + t. Explain how you obtained your answer.
 - (c) If a and b are different positive integers such that $(a^2 4a 21) (b^2 4b 21) = 4$, determine all possible values of a and b. Explain how you obtained your answer.
- 4. In the diagram, four circles of radius 1 with centres P, Q, R, and S are tangent to one another and to the sides of $\triangle ABC$, as shown.



- (a) Determine the size of each of the angles of $\triangle PQS$. Explain how you obtained your answer.
- (b) Determine the length of each side of $\triangle ABC$. Explain how you obtained your answer.
- (c) The radius of the circle with centre R is decreased so that
 - the circle with centre R remains tangent to BC,
 - ullet the circle with centre R remains tangent to the other three circles, and
 - ullet the circle with centre P becomes tangent to the other three circles.

This changes the size and shape of $\triangle ABC$. Determine r, the new radius of the circle with centre R.